

<b>REPORT DOCUMENTATION PAGE</b>				Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503.						
<b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>						
1. REPORT DATE (DD-MM-YYYY) 01-03-2001		2. REPORT DATE FINAL		3. DATES COVERED (From - To) August 1997 - July 2000		
4. TITLE AND SUBTITLE Origins of Nepheloid Layers on Continental Slopes: Laboratory and Field Experimentation				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER N00014-97-1-1011		
				5c. PROGRAM ELEMENT NUMBER		
				5d. PROJECT NUMBER .		
6. AUTHOR(S)  Richard W. Sternberg Erika E. McPhee				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
				8. PERFORMING ORGANIZATION REPORT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) School of Oceanography University of Washington Box 357940 Seattle, WA 98195-7940				10. SPONSOR/MONITOR'S ACRONYM(S)		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5660				11. SPONSORING/MONITORING AGENCY REPORT NUMBER		
12. DISTRIBUTION AVAILABILITY STATEMENT  There are no restrictions to distribution.						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT This was a two-part study to investigate mechanisms for generating intermediate nepheloid layers (INLs) on continental slopes. The first part was a laboratory experiment that generated internal waves on a sloping bed in a stratified fluid which produced INLs. The second part applied the physical understanding gained from the laboratory results to analyze field data collected from the Northern California continental slope as part of the STRATAFORM project. Field data showed that INLs were often associated with regions of critical slope topography where internal wave reflection is common, suggesting that internal waves may be responsible for the generation of slope INLs.						
15. SUBJECT TERMS  continental slopes, internal waves, intermediate nepheloid layers						
16. SECURITY CLASSIFICATION OF: a. REPORT U			17. LIMITATION OF ABSTRACT  UU		18. NUMBER OF PAGES  2	
b. ABSTRACT U			c. THIS PAGE U		19a. NAME OF RESPONSIBLE PERSON Richard Sternberg	
					19b. TELEPHONE NUMBER (Include area code) 206-543-0589	

20010320 044

# ORIGINS OF NEPHELOID LAYERS ON CONTINENTAL SLOPES: LABORATORY AND FIELD EXPERIMENTATION

Richard W. Sternberg

Erika E. McPhee

(Dissertation Advisor: A.R.M. Nowell)

School of Oceanography, Box 357940

University of Washington

Seattle, WA 98195-7940

phone: (206) 543-0589 fax: (206) 543-0275 email: rws@ocean.washington.edu

<http://www.ocean.washington.edu>

Award #N00014-97-1-1011

## RESULTS

When internal waves reflect from a sloping boundary, turbulence and mixing are strongest when the angle of the internal-wave beam is the same as that of the sloping boundary. In the laboratory, mixing of stratified fluid causes intrusions to spread away from the boundary. It is proposed that this process might occur along ocean margins and could be the mechanism by which intermediate nepheloid layers detach and spread seaward from a continental slope.

Laboratory experiments were used to investigate the growth of intrusions due to internal waves reflecting from a sloping boundary. When normalized by the incident energy-density-flux, the average intrusion velocity was found to be a linear function of the parameter  $\omega/\omega_c$ . Evenly spaced layers, indicating thin perturbations in the background density gradient, developed within the mixing region and spread into the tank interior. The vertical spacing between these layers bore a linear relationship to  $\omega/\omega_c$ , and a linear model of internal-wave reflection suggests that these layers may be related to an isopycnal displacement, overturn, scale. Intrusion growth occurred at a range around the critical frequency, and was strongest at slightly supercritical ( $\omega/\omega_c > 1$ ) conditions. A balance between the divergence of energy-density-flux and change in potential energy was suggested to describe the growth of intrusions. Fitting the laboratory results to this theoretical prediction suggested a low net buoyancy flux. The equation for intrusion growth can be used to predict spreading rates for INLs at ocean margins. CTD and transmissometer surveys over the northern California margin showed that INLs could be classified as either shelf INLs, generated between 60 and 200-m depth, or slope INLs, which detached from the continental slope at depths greater than 200 meters. Both shelf and slope INLs were often associated with regions of critical topography. Shelf INLs were strongest in winter, suggesting that their generation is tied to the supply of sediment from winter wave resuspension or Eel River floods. Mooring data suggest that internal-wave reflection is common on this continental slope. Internal-wave reflection occurred in intermittent pulses and may be responsible for the generation of slope INLs at the Eel River margin.

## **PUBLICATIONS**

McPhee, E.E. and D.A. Cacchione (1998) Regional characteristics of suspended sediments at intermediate slope depths on the Northern California continental slope. Talk presented at AGU Fall Meeting, San Francisco, CA.

McPhee, E.E. (2000) Laboratory experiments on internal-wave generation of boundary-layer intrusions and nepheloid layer detachment. Talk presented at AGU/ASLO Ocean Sciences Meeting, San Antonio, TX.

McPhee, E.E. (2000) Internal-wave mixing along sloping boundaries: A mechanism for generating intermediate nepheloid layers. Doctoral dissertation, University of Washington, Seattle, WA.